arXiv:1208.0834v1 [hep-ph] 3 Aug 2012

Bonn-TH-2012-19
IFIC/12-57
CFTP/12-012

A novel way of constraining WIMPs annihilations in the Sun: MeV neutrinos

Nicolás Bernal[1], Justo Martín-Albo[2] and Sergio Palomares-Ruiz[3]

[1]*Bethe Center for Theoretical Physics and Physikalisches Institut,*
Universität Bonn, Nußallee 12, D-53115 Bonn, Germany

[2]*Instituto de Física Corpuscular (IFIC), CSIC-Universitat de València,*
Apartado de Correos 22085, E-46071 Valencia, Spain and

[3]*Centro de Física Teórica de Partículas (CFTP),*
Instituto Superior Técnico, Universidade Técnica de Lisboa,
Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal

Abstract

Annihilation of dark matter particles accumulated in the Sun would produce a flux of high-energy neutrinos whose prospects of detection in neutrino telescopes and detectors have been extensively discussed in the literature. However, for many channels into Standard Model particles, there would also be a flux of neutrinos in the MeV range from the decays at rest of muons and positively charged pions. These low-energy neutrinos have never been considered before and they open the possibility to also constrain dark matter annihilation in the Sun into e^+e^-, $\mu^+\mu^-$ or light quarks. Here we perform a detailed analysis using the recent Super-Kamiokande data in the few tens of MeV range to set limits on the WIMP-nucleon scattering cross section for different annihilation channels.

Introduction.—

There is overwhelming evidence of the existence of a massive non-baryonic dark component which contributes to about 80% of the energy budget of the Universe [1], being a weakly interacting massive particle (WIMP), with mass lying from the GeV to the TeV scale, one of the most popular candidates.

One of the different proposed strategies to detect WIMPs is to search for the flux of high-energy neutrinos from the annihilations of WIMPs accumulated in the center of the Sun [2]. Many different studies have evaluated the prospects of detection of these neutrinos with neutrino telescopes/detectors [3]. However, previous works have focused on WIMPs annihilations into hadronic or $\tau^+\tau^-$ channels. On the other hand, annihilations into $\mu^+\mu^-$ or light quarks have always been neglected as muons and pions lose energy very effectively in the dense regions where they would be produced and then would decay at rest, giving rise to neutrinos in the MeV range. Likewise, annihilations into e^+e^- have never been considered, as e^+/e^- would not produce directly neutrinos. Nevertheless, in their propagation they could interact with nuclei and produce pions, which would be stopped. The π^- would then get captured and subsequently absorbed by the nuclei of the medium without decaying, but the π^+ would decay at rest, producing as well a flux of MeV neutrinos. On the other hand, hadronic and $\tau^+\tau^-$ channels, along with heavy mesons (the source of the high-energy neutrinos considered so far), would also produce pions. The energies of these neutrinos lie at the energy range where the diffuse supernova neutrino background (DSNB) is searched for by detectors as Super-Kamiokande (SK).

Here we consider, for the first time, the potential signal of these low-energy neutrinos from WIMPs annihilations in the Sun and use the most recent SK data [4] to set bounds on the scattering cross section of WIMPs off nucleons for different annihilation channels.

MeV neutrinos from WIMPs annihilations in the Sun.—

Galactic WIMPs could get gravitationally bound within the Sun if after many elastic scatterings off solar nuclei they lose energy and their velocity gets lower than the Sun's escape velocity. Typically, WIMPs would make many orbits between interactions, so their density could be approximated as an isothermal sphere, $n_\chi(r) \propto e^{-m_\chi \phi(r)/T_\chi}$ [5, 6], with $\phi(r)$ the gravitational potential and T_χ the WIMPs temperature, calculated following Ref. [5] and using the Standard Solar Model (SSM) [7].

The evolution of the number of WIMPs in the Sun depends on the capture rate, $C_\odot$, on the annihilation cross section times the relative velocity per unit volume, $A_\odot$, and on the evaporation rate, $E_\odot$, which is only relevant for low-mass WIMPs. We use the capture rates calculated in DarkSUSY [8] and we calculate $A_\odot$ and $E_\odot$ following Ref. [9] and using the SSM [7]. The WIMPs annihilation rate now ($t = t_\odot = 4.57$ Gyr) is given by [9]

$$\Gamma(m_\chi, \sigma_\chi) = \frac{1}{2} C_\odot \left(\frac{\tanh(\kappa\, t_\odot/\tau_E)}{\kappa + \frac{1}{2} E_\odot\, \tau_E\, \tanh(\kappa\, t/\tau_E)} \right)^2 , \tag{1}$$

where $\tau_E = (A_\odot C_\odot)^{-1/2}$ is the equilibration time scale in the absence of evaporation and $\kappa = (1 + (E_\odot \tau_E/2)^2)^{1/2}$. In this work, we assume an annihilation cross section typical of thermal WIMPs, $\langle \sigma_A v \rangle = 3 \cdot 10^{-26}\,\mathrm{cm}^3/\mathrm{s}$ and, for the scattering cross sections under consideration, equilibrium is always reached ($t_\odot >> \tau_E$, $\tanh(\kappa\, t_\odot/\tau_E) \simeq 1$).

Being produced in a very dense medium, among all the final products of WIMPs annihilations, only neutrinos can escape. So far, all previous works have focused on the high-energy neutrino flux resulting from the subsequent hadronization, fragmentation and decay of the final states in heavy quarks, gauge bosons or $\tau^+\tau^-$ channels, and have disregarded annihilations into e^+e^-, $\mu^+\mu^-$ or light quarks because they would only produce (if any) a flux of low-energy neutrinos from pion and muon decay at rest. These MeV neutrinos are the focus of this work.

The propagation of the annihilation products would produce pions that would be stopped and could subsequently decay at rest, giving rise to a monochromatic neutrino spectrum at 29.8 MeV. Practically all π^-, after stopping, are captured in an atomic orbit and get absorbed by the nucleus without decaying. Hence, only neutrinos from π^+ decays would contribute. Let us note that the propagation of high-energy e^-/e^+ would also produce small amounts of pions. In addition, all muons produced in pion decays, in the leptonic decay modes of quarks or taus and in the case of direct annihilations into $\mu^+\mu^-$, are stopped and decay at rest. Thus, in addition to the monochromatic spectrum from π^+ decays at rest, neutrinos from μ^+ and μ^- decaying at rest would also contribute (with a well known spectrum below 52.8 MeV) to the final low-energy neutrino flux.

In order to calculate the relative contributions of each type of neutrino spectrum, we simulate all the particles propagation with GEANT4 [10]. To determine for each WIMP mass the average density and composition of the medium where the products of WIMPs annihilation propagate, we take the SSM [7] and use the WIMP distribution in the Sun.

For the simulations, we proceed as follows. For the case of WIMPs annihilating into a pair of leptons we inject the two leptons with energies equal to the WIMP mass directly into GEANT4 and let them propagate and decay. In the case of WIMPs annihilations into quarks, we use PYTHIA 6.4 [11] to hadronize and fragment the initial quarks and do not let decay any of the final particles that are produced. Then, we inject into GEANT4 the full spectrum of all the produced particles and simulate their propagation in the Sun. Finally, we count the number of π^+, π^-, μ^+ and μ^- that decay at rest, which allow us to compute the initial electron and muon neutrino and antineutrino fluxes at the source.

These neutrinos would then propagate from the Sun to the Earth and would be detected via the charged-current interactions of those arriving at the detector in the electron flavor. In order to calculate the electron neutrino and antineutrino fluxes at the Earth, one should take into account that neutrinos mix. For the energies of interest (above 10 MeV) neutrinos propagate adiabatically in the Sun and at the region where they are produced matter effects are dominant, so ν_e ($\bar{\nu}_e$) exit the Sun as almost purely ν_2 (ν_1) and ν_μ ($\bar{\nu}_\mu$) almost as an equal mixture of ν_1 (ν_2) and ν_3. Hence, the probabilities for neutrinos to arrive at the Earth in the electron flavor are

$$\begin{aligned} P(\nu_\mu \to \nu_e) &= \cos^2\theta_{23}\,\cos^2\theta_{12}\,\cos^2\theta_{13} + \sin^2\theta_{13}\,\cos^2\theta_{13}\,\sin^2\theta_{12}\,(1+\sin^2\theta_{12}) \\ P(\bar{\nu}_\mu \to \bar{\nu}_e) &= \cos^2\theta_{23}\,\sin^2\theta_{12}\,\cos^2\theta_{13} + \sin^2\theta_{13}\,\cos^2\theta_{13}\,\sin^2\theta_{12}\,(1+\cos^2\theta_{12}) \\ P(\nu_e \to \nu_e) &= \sin^2\theta_{12}\,\cos^2\theta_{13} + \sin^4\theta_{13} \\ P(\bar{\nu}_e \to \bar{\nu}_e) &= \cos^2\theta_{12}\,\cos^2\theta_{13} + \sin^4\theta_{13}\ . \end{aligned} \tag{2}$$

The final electron neutrino and antineutrino spectra at the detector are obtained by combining the initial fluxes with the above probabilities (and we have neglected the small correction due to the Earth-matter effect). We use the values of the mixing angles from Ref. [12].

Detection of MeV neutrinos with SK.—

SK is a water Čerenkov detector with a fiducial volume of 22.5 ktons. For energies below 52.83 MeV, the best present data come from the search for the DSNB [4], which is split into three phases: SK-I ($t_I = 1497$ days), SK-II ($t_{II} = 794$ days) and SK-III ($t_{III} = 562$ days). The recent analysis [4] has substantially improved over the previous one [13], with better efficiencies, lower energy thresholds and almost twice as much statistics.

The signal at SK is the detection of the positrons (electrons) produced in $\bar{\nu}_e$ (ν_e) charged-current interactions in the detector below $\sim$ 100 MeV. At these energies, the inverse beta-decay cross section ($\bar{\nu}_e p \to n e^+$) is by two orders of magnitude larger that the $\nu - e$ elastic scattering cross section. Below $\sim$80 MeV, this is the dominant interaction of $\bar{\nu}_e$. However, the interactions ν_e and $\bar{\nu}_e$ off oxygen nuclei give non-negligible contributions and need also to be taken into account.

The low energy threshold in the analysis is determined by the ability to remove the radioactive spallation caused by cosmic-ray muons hitting an oxygen nucleus. An improved spallation cut has allowed to reduce the energy threshold used in the previous analysis down to 16 MeV (17.5 MeV) for SK-I/III (SK-II). In addition to this, a number of other cuts were performed, as noise reduction, fiducial volume, solar angle, incoming event, decay electron, pion, Čerenkov angle and other cuts [4]. The maximum energy in the recent SK analysis is 88 MeV, which is also higher than in the previous one.

In this energy range, the two dominant backgrounds which remain after the cuts are the atmospheric ν_e and $\bar{\nu}_e$ background and, mainly, the Michel positrons (electrons) from the decays of low energy muons, produced by atmospheric $\bar{\nu}_\mu$ (ν_μ) with typical energies of about $\sim$200 MeV, which are below detection threshold, the so-called invisible muons. These muons are slowed down rapidly and subsequently decay, mimicking the signal from ν_e or $\bar{\nu}_e$, but with a spectrum whose shape is very well known.

Analysis.—

In this work, we follow closely the analysis performed in Refs. [14], although updating some aspects following the new SK analysis [4].

We have considered both the interactions of $\bar{\nu}_e$ off free protons and the interactions of ν_e and $\bar{\nu}_e$ off bound nucleons. For the inverse beta-decay reaction we use the full cross section [15] and for the interactions off bound nucleons, we consider a relativistic Fermi gas model [16] with a Fermi surface momentum of 225 MeV and a binding energy of 27 MeV. For each of the three SK phases, we have used the corresponding energy-dependent efficiencies [4] and a Gaussian energy resolution function [17].

In order to obtain the upper limit on the WIMP-nucleon scattering cross section, we use the recent data reported by the SK collaboration [4]. Unlike the SK analysis, we only consider the two backgrounds described above and only take the data in the Čerenkov angle region defined as the signal region [4]. We expect this to affect very little our results because the two extra sources of background considered in the SK analysis are subdominant at the percent level and their energy distribution is very flat, while the normalizations of the potential signal and of each background are left free in our analysis, which are the three parameters we fit (α, β and γ, which represent the total number of signal and background events). We consider 18 4-MeV bins (in the interval 16-88 MeV) and perform an unbinned maximum likelihood fit. As in the SK analysis, we obtain the best fit as the combination of parameters that maximizes the likelihood, defined as

$$\mathcal{L} = e^{-(\alpha+\beta+\gamma)} \prod_{l=1}^{18} \left[(\alpha \cdot A_l) + (\beta \cdot B_l) + (\gamma \cdot C_l)\right]^{N_l} , \tag{3}$$

where the sum l is over all energy bins, N_l is the number of detected events in the lth bin, and A_l, B_l and C_l are the fractions (so that for each case the total is normalized to 1) of the WIMPs annihilation signal, Michel positrons and electrons from muon decay and atmospheric ν_e and $\bar{\nu}_e$ spectra that are in the lth bin, respectively. The fractions A_l are calculated using the $\bar{\nu}_e$ and ν_e low-energy fluxes from WIMP annihilations in the Sun as described above. When computing B_l we have taken into account that 18.4% of the μ^- are stopped in water and the spectrum of their decay electron gets distorted. In the calculation of the fractions B_l and C_l we have used the atmospheric neutrino flux calculation with FLUKA [18].

We have also included the energy-independent efficiency systematic error by modifying the likelihood in the way described in Ref. [4], with a total error different for each of the data-taking phases. The final likelihood is maximized for each SK phase separately, so that it remains as a function of just α, the number of signal events. Finally, the three likelihoods are calculated as a function of the number of signal events/year and multiplied. The 90% confidence level (CL) limit on the number of signal events/year is determined by

$$\frac{\int_0^{\alpha_{90}} \mathcal{L}_{\rm tot}(\alpha)\, d\alpha}{\int_0^{\infty} \mathcal{L}_{\rm tot}(\alpha)\, d\alpha} = 0.9 \ . \tag{4}$$

The limit on the scattering cross section (σ_χ^{90}) is then obtained by solving

$$\Gamma(m_\chi, \sigma_\chi^{90})\, A^{\rm tot} = \alpha_{90} \quad , \; A^{\rm tot} \equiv \frac{\sum_{SK} A^{SK} t_{SK}}{\sum_{SK} t_{SK}} , \tag{5}$$

where A^{SK} is the number of events per WIMPs annihilation for each SK phase at the detector.



Figure 1: ***Limits on the spin-dependent scattering cross section of WIMPs off protons at 90% CL for different annihilation channels.*** *The limits from SIMPLE, PICASSO and CDMS [19] are shown with black lines. The limits from SK searches of GeV neutrinos are depicted for two annihilation channels [20]. The DAMA/LIBRA [21] regions (at 90% CL and 3σ CL) are also shown as interpreted in Refs. [22].*

The results for spin-dependent (off protons) and spin-independent cross sections are shown in Figs. 1 and 2, respectively, along with the bounds from direct detection searches and the analysis of GeV neutrinos at SK. The most stringent bounds we obtain are for WIMPs annihilations into light quarks, which is the channel that generates the largest amount of pions. Whereas for spin-independent, the limits are one or two orders of magnitude weaker than the ones obtained with direct searches, for spin-dependent and low WIMP mass, they are comparable to (or even more stringent than) them. We note that, in this case, the limits for WIMPs annihilations into quarks exclude the DAMA/LIBRA [21] region at 90% CL. We also obtain bounds for the case of WIMPs annihilations into e^+e^-,



Figure 2: ***Limits on the spin-independent scattering cross section of WIMPs at 90% CL for different annihilation channels.*** *The limits from CDMS, COGENT, XENON10 and XENON100 [23] are shown with black lines. The limits from SK searches of GeV neutrinos are depicted for two annihilation channels [20]. The DAMA/LIBRA [21] regions (at 90% CL and* 3σ *CL) are also shown as interpreted in Refs. [22].*

although very weak. The limit for the $\mu^+\mu^-$ channel follows the dependence of the capture rate with the WIMP mass, as the initial muons are always stopped. On the other hand, for the $\tau^+\tau^-$ channel, at low masses, the number of muons decaying at rest is smaller because a fraction of taus goes to electrons and all the produced π^- get absorbed without decaying. However, the passage through the solar medium of the products of tau decay gives rise to a number of pions that increases with energy and hence, with the WIMP mass (the same trend is seen for the case of WIMPs annihilations into quarks). This explains why these limits cross each other.

Conclusions.—

The potential signal of GeV neutrinos produced after WIMPs annihilations in the Sun has been extensively studied so far [3]. However, these searches do not consider annihilations into light quarks, $\mu^+\mu^-$ or e^+e^-, as these channels do not produce GeV neutrinos. However, the propagation in the Sun of the products of WIMPs annihilations into quarks or leptons would always produce pions and muons that after being stopped would decay at rest. In this work, we have considered these MeV neutrinos from pion and muon decay at rest, which represents a novel way of constraining WIMPs annihilations in the Sun. In order to do so, we have used the SK data on the DSNB [4] and have performed and analogous analysis. Our results, Figs. 1 and 2, show that these new limits are complementary to those from direct searches, mainly for low-mass WIMPs.

Note added: As this work was being completed, we learned of an independent study with similar results by C. Rott, J. Siegal-Gaskins and J. F. Beacom [24] that will be submitted to arXiv simultaneously.

Acknowledgments.—

We thank K. Bays and C. Peña-Garay for providing us with useful information, M. Vicente-Vacas for discussions and M. Sorel for encouragement. SPR thanks the Galileo Galilei Institute for Theoretical Physics, where parts of this work were done, for its hospitality. NB is supported by the DFG TRR33 'The Dark Universe'. JMA is supported by the Spanish Grant CONSOLIDER-Ingenio 2010 CSD2008-0037 (CUP) of the MINECO. SPR is partially supported by the Portuguese FCT through CERN/FP/123580/2011 and CFTP-FCT UNIT 777, which are partially funded through POCTI (FEDER) and by the Spanish Grant FPA2011-23596 of the MINECO.

---

[1] G. Bertone, D. Hooper and J. Silk, Phys. Rept. **405**, 279 (2005) [arXiv:hep-ph/0404175].

[2] J. Silk, K. A. Olive and M. Srednicki, Phys. Rev. Lett. **55**, 257 (1985); M. Srednicki, K. A. Olive and J. Silk, Nucl. Phys. B **279**, 804 (1987); T. K. Gaisser, G. Steigman and S. Tilav, Phys.

Rev. D **34**, 2206 (1986).

[3] M. Kamionkowski, Phys. Rev. D **44**, 3021 (1991); A. Bottino *et al.*, Phys. Lett. B **265**, 57 (1991); F. Halzen, T. Stelzer and M. Kamionkowski, Phys. Rev. D **45**, 4439 (1992); L. Bergstrom, J. Edsjo and P. Gondolo, Phys. Rev. D **55**, 1765 (1997) [hep-ph/9607237]; V. D. Barger *et al.*, Phys. Rev. D **65**, 075022 (2002) [hep-ph/0105182]; M. Cirelli *et al.*, and F. Vissani, Nucl. Phys. B **727**, 99 (2005) [Erratum-ibid. B **790**, 338 (2008)] [hep-ph/0506298]; O. Mena, S. Palomares-Ruiz and S. Pascoli, Phys. Lett. B **664**, 92 (2008) [arXiv:0706.3909 [hep-ph]]; C. R. Das *et al.*, arXiv:1110.5095 [hep-ph].

[4] K. Bays *et al.* [Super-Kamiokande Collaboration], Phys. Rev. D **85**, 052007 (2012) [arXiv:1111.5031 [hep-ex]].

[5] D. N. Spergel and W. H. Press, Astrophys. J. **294**, 663 (1985).

[6] J. Faulkner and R. L. Gilliland, Astrophys. J. **299**, 994 (1985).

[7] A. M. Serenelli, W. C. Haxton and C. Pena-Garay, Astrophys. J. **743**, 24 (2011) [arXiv:1104.1639 [astro-ph.SR]].

[8] P. Gondolo *et al.*, JCAP **0407**, 008 (2004) [astro-ph/0406204].

[9] K. Griest and D. Seckel, Nucl. Phys. B **283**, 681 (1987) [Erratum-ibid. B **296**, 1034 (1988)].

[10] S. Agostinelli *et al.* [GEANT4 Collaboration], Nucl. Instrum. Meth. A **506**, 250 (2003); J. Allison *et al.*, IEEE Trans. Nucl. Sci. **53**, 270 (2006).

[11] T. Sjostrand, S. Mrenna and P. Z. Skands, JHEP **0605**, 026 (2006) [hep-ph/0603175].

[12] D. V. Forero, M. Tortola and J. W. F. Valle, arXiv:1205.4018 [hep-ph].

[13] M. Malek *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **90**, 061101 (2003) [hep-ex/0209028].

[14] S. Palomares-Ruiz and S. Pascoli, Phys. Rev. D **77**, 025025 (2008) [arXiv:0710.5420 [astro-ph]]; S. Palomares-Ruiz, Phys. Lett. B **665**, 50 (2008) [arXiv:0712.1937 [astro-ph]].

[15] A. Strumia and F. Vissani, Phys. Lett. B **564**, 42 (2003) [astro-ph/0302055]; P. Vogel and J. F. Beacom, Phys. Rev. D **60**, 053003 (1999) [hep-ph/9903554].

[16] R. A. Smith and E. J. Moniz, Nucl. Phys. B **43**, 605 (1972) [Erratum-ibid. B **101**, 547 (1975)].

[17] J. Hosaka *et al.* [Super-Kamiokande Collaboration], Phys. Rev. D **73**, 112001 (2006) [hep-ex/0508053]; J. P. Cravens *et al.* [Super-Kamiokande Collaboration], Phys. Rev. D **78**, 032002 (2008) [arXiv:0803.4312 [hep-ex]]; K. Abe *et al.* [Super-Kamiokande Collaboration], Phys. Rev. D **83**, 052010 (2011) [arXiv:1010.0118 [hep-ex]].

[18] G. Battistoni *et al.*, Astropart. Phys. **19**, 269 (2003) [Erratum-ibid. **19**, 291 (2003)] [hep-ph/0207035]; and Astropart. Phys. **23**, 526 (2005).

[19] M. Felizardo *et al.* [SIMPLE Collaboration], Phys. Rev. Lett. **108**, 201302 (2012) [arXiv:1106.3014 [astro-ph.CO]]; S. Archambault *et al.* [PICASSO Collaboration], Phys. Lett. B **711**, 153 (2012) [arXiv:1202.1240 [hep-ex]]; E. Behnke *et al.* [COUPP Collaboration], arXiv:1204.3094 [astro-ph.CO].

[20] T. Tanaka *et al.* [Super-Kamiokande Collaboration], Astrophys. J. **742**, 78 (2011) [arXiv:1108.3384 [astro-ph.HE]]; R. Kappl and M. W. Winkler, Nucl. Phys. B **850**, 505 (2011) [arXiv:1104.0679 [hep-ph]].

[21] R. Bernabei *et al.* [DAMA/LIBRA Collaboration], Eur. Phys. J. C **56**, 333 (2008) [arXiv:0804.2741 [astro-ph]]; and Eur. Phys. J. C **67**, 39 (2010) [arXiv:1002.1028 [astro-ph.GA]].

[22] C. Savage *et al.*, JCAP **0904**, 010 (2009) [arXiv:0808.3607 [astro-ph]]; and Phys. Rev. D **83**, 055002 (2011) [arXiv:1006.0972 [astro-ph.CO]].

[23] C. E. Aalseth *et al.* [CoGeNT Collaboration], Phys. Rev. Lett. **106**, 131301 (2011) [arXiv:1002.4703 [astro-ph.CO]]; D. S. Akerib *et al.* [CDMS Collaboration], Phys. Rev. D **82**, 122004 (2010) [arXiv:1010.4290 [astro-ph.CO]]; J. Angle *et al.* [XENON10 Collaboration], Phys. Rev. Lett. **107**, 051301 (2011) [arXiv:1104.3088 [astro-ph.CO]]; E. Aprile *et al.* [XENON100 Collaboration], arXiv:1207.5988 [astro-ph.CO].

[24] C. Rott, J. Siegal-Gaskins and J. F. Beacom, submitted to arXiv.